

ALL ORDERS MUST BE SIGNED AND DATED BY THE REFERRING PROVIDER

ALLERGIES/REACTIONS (REQUIRED):

Yakima Outpatient Infusion Care
808 N 39th Ave Yakima WA 98902
Phone: 509-575-1174
Fax: 509-577-5021

ORDERS WITH CHECK BOXES

When an order is optional (those with check boxes), providers are responsible for indicating a check mark in the box next to the order. Orders left unchecked will not be initiated.

CODE STATUS

Patients will be considered FULL CODE unless marked otherwise. If the patient has a POLST, advance directive, or living will, please include a copy with the orders.

BELIMUMAB (Benlysta)

Patient Name: _____ Requested Start Date: ____/____/____
Date of Birth: ____/____/____ Patient Weight: _____ kg Patient Height: _____

DIAGNOSIS & ICD-10 CODE:

- Systemic Lupus Erythematosus (ICD-10: _____)
- Other: _____ (ICD-10: _____)

REQUIRED: H&P with documentation to support above diagnosis including ICD-10 code and supporting labs

****If required documentation not received with this order, scheduling of treatment will be delayed until complete information is available****

ROUTINE LABS: CMP | CBC w/ diff | ANC | LFT | CRP | Lipid Panel | ESR | Other: _____

ROUTINE LAB FREQUENCY: Each Infusion | Annually | Other: _____

ACCESS: Access and maintain IV site or Port-A-Cath in accordance with the appropriate MYM OIC P&Ps

TREATMENT REGIMEN: (another brand of drug, identical in form and content may be dispensed unless "DAW" or "BRAND ONLY" is written next to the drug name)

PRE-MEDICATIONS (OPTIONAL) – Give 30 minutes prior to infusion

- Acetaminophen 650 mg PO x1 Diphenhydramine 25 mg PO x1
- Loratadine 10 mg PO x1
- Other: _____

BELIMUMAB (Benlysta) in 250 mL 0.9% Normal Saline and infused over 60 minutes. (dose may be rounded to nearest vial size PRN)

- INDUCTION**
10 mg/kg = _____ mg IV every 2 weeks x 3 doses then every 4 weeks thereafter
- MAINTENANCE**
10 mg/kg = _____ mg IV every 4 weeks

MONITORING: Vitals at baseline and at completion of infusion.

SUPPORTIVE CARE: Administer hypersensitivity reaction/anaphylaxis management per MYMH OIC protocol as necessary.

DISCHARGE: 30 minutes after infusion is complete when vital signs are stable and hypersensitivity symptoms are absent. Waiting period can be waived by patient on subsequent infusions.

Provider Signature: _____ Date: _____
Print name: _____ Phone # _____ Fax # _____

- NEW REFERRAL UPDATED REFERRAL

****Expires 12 months from written date****

Patient Identification - Attach Patient Label

Name: _____
MRN: _____
Age / Sex and Gender: _____

BELIMUMAB (Benlysta)
MultiCare 
Yakima Memorial Hospital