

ALL ORDERS MUST BE SIGNED AND DATED BY THE REFERRING PROVIDER

ALLERGIES/REACTIONS (REQUIRED):

Yakima Outpatient Infusion Care
808 N 39th Ave Yakima WA 98902
Phone: 509-575-1174
Fax: 509-577-5021

ORDERS WITH CHECK BOXES

When an order is optional (those with check boxes), providers are responsible for indicating a check mark in the box next to the order. Orders left unchecked will not be initiated.

CODE STATUS

Patients will be considered FULL CODE unless marked otherwise. If the patient has a POLST, advance directive, or living will, please include a copy with the orders.

RISANKIZUMAB-RZAA (Skyrizi) INDUCTION

Patient Name: _____ Requested Start Date: ____/____/____
Date of Birth: ____/____/____ Patient Weight: _____ kg Patient Height: _____

DIAGNOSIS & ICD-10 CODE:

- Crohn's disease (ICD-10: _____)
- Other: _____ (ICD-10: _____)

REQUIRED: H&P with documentation to support above diagnosis including ICD-10 code and supporting labs
****If required documentation not received with this order, scheduling of treatment will be delayed until complete information is available****

REQUIRED BASELINE LABS & INFORMATION:

- ✓ CBC & CMP within 30 days prior to first infusion
- ✓ Negative Latent TB Test (Date: _____ | QuantiFERON Gold | PPD | Chest X-Ray | Other: _____)

ACCESS: Access and maintain IV site or Port-A-Cath in accordance with the appropriate MYM OIC P&Ps

TREATMENT REGIMEN: (another brand of drug, identical in form and content may be dispensed unless "DAW" or "BRAND ONLY" is written next to the drug name)

Risankizumab-rzaa (Skyrizi) IV Induction

Subcutaneous maintenance therapy to start at week 12 after induction dose – Please arrange with the patient's preferred outpatient or specialty pharmacy.

- ✓ Induction: 600 mg IV infused over at least 1 hour at week 0, week 4, and week 8

MONITORING: Vitals at baseline and at completion of infusion.

SUPPORTIVE CARE: Administer hypersensitivity reaction/anaphylaxis management per MYMH OIC protocol as necessary.


DISCHARGE: 30 minutes after infusion is complete when vital signs are stable and hypersensitivity symptoms are absent. Waiting period can be waived by patient on subsequent infusions.

Provider Signature: _____ Date: _____
Print name: _____ Phone # _____ Fax # _____

****Expires 12 months from written date****

Patient Identification - Attach Patient Label

Name:
MRN:
Age / Sex and Gender:

RISANKIZUMAB (Skyrizi) INDUCTION
MultiCare 
Yakima Memorial Hospital